



## Background Information on Edible Insects & Human Health

Written By: Valerie Stull, MPH

Edible insects have been consumed throughout human history<sup>1,2</sup> across Asia, the Americas, Africa,<sup>3</sup> and Australia.<sup>4</sup> Even native populations of North America, especially in the west were known to consume insects and expend significant effort to harvest them.<sup>3</sup> Insects served as an important component of the diet for many populations, traditionally making up more than 50% of animal protein intake in parts of central Africa<sup>5</sup> and containing macronutrient values similar to that of wild meat, shellfish, nuts, pulses, and some vegetables.<sup>6</sup> Entomophagy—the practice of eating insects—continues today, as nearly 2 billion people<sup>7</sup> from 3,000 ethnic groups in 130 countries<sup>8</sup> regularly consume insects, selecting from more than 1900 known edible species.<sup>9</sup> These people are spread across 80% of the world's populations.<sup>10</sup> In Africa alone, at least 500 species are consumed in 40 different nations.<sup>11</sup> Even within the last century, some groups have relied heavily on insects, such as residents of the Kwango District in the Democratic Republic of Congo who garner nearly a third of their animal protein intake from insects.<sup>12,13</sup> Today, there are more than 24 food companies selling insect-based food products in North America alone, and the edible insect industry is growing rapidly in Europe as well. In 2015, the European Food Safety Authority took preliminary steps to assess the food safety of edible insects, as they are widely available in Europe thanks to trade with Asian producers and the emergence of new businesses in Europe. The Food and Agriculture Organization of the United Nations (FAO) has deemed many edible insects safe to eat when properly prepared.<sup>7</sup>

Edible insects are nutritionally dense, offering a good source of bioavailable animal protein for those that partake.<sup>14</sup> Many edible species contain high crude protein levels,<sup>15,16</sup> with between 40 and 75% protein by dry weight on average,<sup>17</sup> higher than that of ground beef.<sup>18</sup> Insect protein quality is also superior, as the amino acid content of most edible insects is comparable to WHO recommendations for amino acids required for human nutrition,<sup>19</sup> and insects commonly contain all essential amino acids.<sup>20</sup> In general, insects are rich in fatty acids,<sup>15</sup> with numerous species supplying ample poly-unsaturated fatty acids (including essential linoleic and linolenic acids).<sup>21</sup> The exoskeleton of insects contains chitin, making insects a source of dietary fiber, with chitin accounting for about 10% of whole dried insects.<sup>15</sup> Chitin, a modified polysaccharide, is abundant in nature and has been evaluated for its health-promoting properties. For example, chitin-glucan derived from *Aspergillus niger* mycelium reduced oxidized low-density lipoprotein in adults, a risk factor for atherosclerosis, when consumed at a rate of 4.5g/day.<sup>22</sup> Insects are also typically good sources of minerals including potassium, calcium, copper, magnesium,<sup>17</sup> manganese, phosphorus, and selenium,<sup>23-24</sup> as well as the ever important iron and zinc,<sup>25</sup> which are crucial for growth. Some insect types are particularly high in B vitamins,<sup>17</sup> such as biotin, riboflavin, pantothenic acid, and folate.<sup>23</sup> Caterpillars, for example, contain as much as 300 times more than daily requirement for adults of iron, along with ample zinc.<sup>26</sup> As such, insect consumption has been proposed as one way to decrease iron and zinc deficiency in developing countries<sup>25</sup> and reduce malnutrition globally. It should be noted, however, that the nutritional value of insects can vary greatly by species, lifecycle,<sup>27</sup> and feed source(s).<sup>28</sup>

From an environmental perspective, insects are a plentiful natural resource, comprising ~1 million of the 1.4 million species on the planet.<sup>7</sup> The environmental impact of edible insects is significantly lower than traditional livestock,<sup>7,20,29</sup> as they need less land, water, and feed to survive and thrive,<sup>7</sup> and emit fewer greenhouse gases (GHGs).<sup>29</sup> Their high feed-conversion ratio,<sup>20</sup> ectothermic thermoregulation, and large edible body mass percentage<sup>30</sup> contribute to overall sustainability and desirability. Production of 1 kilogram (kg) of live animal mass requires about 2.5kg of feed of poultry, 5kg for swine, and 10kg for beef,<sup>31</sup> whereas crickets demand as little as 1.7 kg feed per kg of mass gained.<sup>20</sup> Excluding scarab beetles, cockroaches, and termites, edible insects do not produce methane (CH<sub>4</sub>),<sup>32</sup> an especially powerful GHG with a significantly higher global warming potential (GWP) than carbon dioxide (CO<sub>2</sub>).<sup>33</sup> The total GHG emissions of several common edible insects (crickets, mealworms, and locusts) are lower by a factor of about 100 than pigs or beef cattle.<sup>29</sup>

Despite the numerous perks and overall global popularity of edible insects, a more research is needed to explore the health impact of entomophagy on consumers, including their ability to curb protein deficiencies or undernutrition, influence the microbiota of humans, and address micronutrient deficiencies. Research exploring and addressing challenges with current collection, processing, and consumption methods, as well as new innovations that harness the relatively untapped potential of edible insects are needed.

## References:

1. Bodenheimer FS. *Insects as Human Food: A Chapter of the Ecology of Man*. W. Junk; 1951.
2. Defoliart GR. Edible insects as minilivestock. *Biodivers Conserv*. 1995;4(3):306-321. doi:10.1007/BF00055976.
3. Sutton MQ. *Insects As Food: Aboriginal Entomophagy in the Great Basin*. Menlo Park, CA: Ballena Pr; 1988.
4. Yen AL. Insect and Other Invertebrate Foods of the Australian Aborigines. In: Paoletti MD, ed. *Ecological Implications of Minilivestock. Potential of Insects, Rodents, Frogs and Snails*. Enfield, (NH): Science Publishers; 2005:367-387.
5. Paoletti MG. *Ecological Implications of Minilivestock Potential of Insects, Rodents, Frogs, and Snails*. Enfield, (NH): Science Publishers; 2005. <http://site.ebrary.com/id/10257792>. Accessed August 13, 2015.
6. Raubenheimer D, Rothman JM, Pontzer H, Simpson SJ. Macronutrient contributions of insects to the diets of hunter-gatherers: A geometric analysis. *Journal of Human Evolution*. 2014;71:70-76. doi:10.1016/j.jhevol.2014.02.007.
7. van Huis A, Van Itterbeeck J, Klunder H, et al. *Edible Insects Future Prospects for Food and Feed Security*. Rome: Food and Agriculture Organization of the United Nations (FAO); 2013. <http://www.fao.org/docrep/018/i3253e/i3253e00.htm>. Accessed November 13, 2013.
8. Ramos-Elorduy J. Anthro-entomophagy: Cultures, evolution and sustainability. *Entomological Research*. 2009;39(5):271-288. doi:10.1111/j.1748-5967.2009.00238.x.
9. van Huis A, Van Itterbeeck J, Klunder H, et al. *Edible Insects: Future Prospects for Food and Feed Security*. Rome: Food and Agriculture Organization of the United Nations (FAO); 2013. <http://www.fao.org/docrep/018/i3253e/i3253e00.htm>. Accessed November 13, 2013.
10. Srivastava S, Babu N, Pandey H. Traditional insect bioprospecting - As human food and medicine. *Indian J Tradit Knowl*. 2009;8(4):485-494.
11. Jongema Y. List of edible insects of the world (April 1, 2014). Wageningen UR. <http://www.wageningenur.nl/en/Expertise-Services/Chair-groups/Plant-Sciences/Laboratory-of-Entomology/Edible-insects/Worldwide-species-list.htm>. Published 2014. Accessed December 13, 2014.
12. Gomez PA, Halut R, Collin A. Production de proteines animales au Congo. *Bull Agric Congo*. 1961.
13. DeFoliart GR. Insects as Food: Why the Western Attitude Is Important. *Annual Review of Entomology*. 1999;44(1):21.
14. Verkerk MC, Tramper J, van Trijp JCM, Martens DE. Insect cells for human food. *Biotechnology Advances*. 2007;25(2):198-202. doi:10.1016/j.biotechadv.2006.11.004.
15. Belluco S, Losasso C, Maggiolini M, Alonzi CC, Paoletti MG, Ricci A. Edible Insects in a Food Safety and Nutritional Perspective: A Critical Review. *Comprehensive Reviews in Food Science and Food Safety*. 2013;12(3):296-313. doi:10.1111/1541-4337.12014.
16. Melo V, Garcia M, Sandoval H, Jiménez HD, Calvo C. Quality proteins from edible indigenous insect food of Latin America and Asia. *Emirates Journal of Food and Agriculture*. 2011;23(3):283-289.
17. Schabel HG. Forest insects as food: a global review. In: Food and Agriculture Organization of the United Nations (FAO); 2010:37-64.
18. Gahukar RT. Entomophagy and human food security. *International Journal of Tropical Insect Science*. 2011;31(03):129-144. doi:10.1017/S1742758411000257.
19. Rumpold BA, Schlüter OK. Potential and challenges of insects as an innovative source for food and feed production. *Innovative Food Science & Emerging Technologies*. 2013;17:1-11. doi:10.1016/j.ifset.2012.11.005.
20. Collavo A, Glew RH, Huang YS, Chuang LT, Bosse R, Paoletti MG. House cricket small-scale farming. In: Paoletti MG, ed. *Ecological Implications of Minilivestock: Potential of Insects, Rodents, Frogs and Snails*. Enfield, (NH): CRC Press; 2005:519-544.
21. Womehi HM, Linder M, Tiencheu B, et al. Oils of insects and larvae consumed in Africa: potential sources of polyunsaturated fatty acids. *Oléagineux, Corps gras, Lipides*. 2009;16(4-5-6):230-235. doi:10.1051/ocl.2009.0279.
22. Bays HE, Evans JL, Maki KC, et al. Chitin-glucan fiber effects on oxidized low-density lipoprotein: a randomized controlled trial. *Eur J Clin Nutr*. 2013;67(1):2-7. doi:10.1038/ejcn.2012.121.
23. Rumpold BA, Schlüter OK. Nutritional composition and safety aspects of edible insects. *Mol Nutr Food Res*. 2013;57(5):802-823. doi:10.1002/mnfr.201200735.
24. Finke MD. Complete nutrient composition of commercially raised invertebrates used as food for insectivores. *Zoo Biol*. 2002;21(3):269-285. doi:10.1002/zoo.10031.
25. Christensen DL, Orech FO, Mungai MN, Larsen T, Friis H, Aagaard-Hansen J. Entomophagy among the Luo of Kenya: a potential mineral source? *International Journal of Food Sciences & Nutrition*. 2006;57(3/4):198-203.
26. DeFoliart GR. Insects as human food: Gene DeFoliart discusses some nutritional and economic aspects. *Crop Protection*. 1992;11(5):395-399. doi:10.1016/0261-2194(92)90020-6.
27. Nowak V, Persijn D, Rittenschöber D, Charrondière UR. Review of food composition data for edible insects. *Food Chemistry*. doi:10.1016/j.foodchem.2014.10.114.
28. Oonincx DGAB, van Broekhoven S, van Huis A, van Loon JJA. Feed Conversion, Survival and Development, and Composition of Four Insect Species on Diets Composed of Food By-Products. *PLoS ONE*. 2015;10(12):1-20. doi:10.1371/journal.pone.0144601.
29. Oonincx DGAB, van Itterbeeck J, Heetkamp MJW, van den Brand H, van Loon JJA, van Huis A. An Exploration on Greenhouse Gas and Ammonia Production by Insect Species Suitable for Animal or Human Consumption. *PLoS One*. 2010;5(12). doi:10.1371/journal.pone.0014445.
30. Nakagaki BJ, DeFoliart GR. Comparison of Diets for Mass-Rearing Acheta domesticus (Orthoptera: Gryllidae) as a Novelty Food, and Comparison of Food Conversion Efficiency with Values Reported for Livestock. *Journal of Economic Entomology*. 1991;84(3):891-896.
31. Smil V. Worldwide transformation of diets, burdens of meat production and opportunities for novel food proteins. *Enzyme and Microbial Technology*. 2002;30(3):305-311. doi:10.1016/S0141-0229(01)00504-X.
32. Hackstein JH, Stumm CK. Methane production in terrestrial arthropods. *Proc Natl Acad Sci U S A*. 1994;91(12):5441-5445.
33. Kemp DD. *Exploring Environmental Issues: An Integrated Approach*. Routledge; 2004.